The Bryologist 91(4), 1988, pp. 385–396
Copyright © 1988 by the American Bryological and Lichenological Society, Inc.

Index to Volume 91 of THE BRYOLOGIST[1,2]

Acanthocladium alboalare, *37*; bifarium, *37*; polymorphum, *37*; scabrifolium, *37*; tanytrichum, *37*; tenuisetum, 37

Acanthorrhynchium papillatum, 34+; papillatum fo. penangense, 34

Acarospora fuscata, 309+; molybdina, 309+; smaragdula, 309+

Acaulon muticum var. rufescens, 214+

Acid rain, simulated, effects on Tomenthypnum nitens and Scorpidium scorpioides, 121

Acroporium aciphyllum, 34; albidissimum, 34; asperifolium, 34; bogoricum, *37*; braunii, *37*; brevipes, 34; clastobryelloides, *37*; complanatum, *37*; condensatum, 34; convolutum, 34; decipiens, 35+; denticulatum, 35; cf. dicranoides, 35; diminutum, 35+; divaricatum, 37; falcifolium, 35; hamulatum, 35; hamulatum fo. procumbens, 35; hermaphroditum, 35+; hyalinum, 35+; laevibogoricum, *37*; lamprophyllum, 35; leucophyllum, 35; longicuspis, 35+; malayanum, *37*; monoicum, *37*; obscurum, 35+; oxyporum, 35; perangustifolium, 37; procerum, 35; punctuliferum, 35; pungens, 338; pycnophyllum, *37*; ridleyi, 35; rigens, 35+; rigens var. dicranolomoides, 35+; ruficaule, *37*+; rufum, 35+; secundum, 35+; secundum var. angustifolium, 35; secundum var. latifolium, 35; secundum var. longisetum, 35; secundum var. minus, 35; serrulatum, *37*; sigmatodontium, 35; stramineum, 35, *37*+; subulatum, *37*+; surculare, *37*; turgidum, 35; cf. vincensianum, 35

Actinodontium adscendens, 33; portoricense, new to Colombia, 130

Adelothecium bogotense, 337

Aerobryidium aureo-nitens, 32; crispifolium, 32; filamentosum, 32

Aerobryopsis filamentosum, 37; leptosigmata, 32; longissima, *37*; longissima var. condensata, *37*; punctulatum, *37*; subleptostigmata, 32; wallichii, 32+

Agyrium rufum, 309+

Alaska, Staurothele discedens, Gyalidea lecideopsis var. convarians and Polyblastia cucurbitula in, 86

Alectoria sarmentosa, 309+

ALLEN, B. H. A contribution to the moss flora of Peru, including a new species of Dicranum, 91

ALLEN, B. H. *See* PURSELL, R. A., M. A. BRUGGEMAN-NANNENGA & B. H. ALLEN, 202

ALLEN, B. H. (review). J.-P. Frahm, a revised list of the Campylopus species of the world, 158

ALLEN, B. H. (review). D. G. Long, Illustrated Moss Flora of Arctic North America and Greenland. 1. Polytrichaceae, 252

Amblystegium fluviatile, 98; noterophyllum, 98; riparium, 98; serpens var. juratzkanum, 98; serpens var. serpens, 98; tenax var. spinifolium, 98; tenax var. tenax, 98; trichopodium, 98; varium, 98

Amphidium mougeotii, 98

Amygdalaria panaeola, 309+

Anacamptodon splachnoides, 98

Anacolia laevisphaera, 336

Anaptychia palmulata, 309+

ANDERSON, L. E. & W. R. BUCK. Entodon hampeanus C. Müll., new and extinct in the United States, 113

Andreaea acutifolia, new to Colombia, 334+; blyttii, 358; brevipes, 335; nivalis, 358; rothii, 98

Andreaeobryum macrosporum, 88

ANDRUS, R. E. Validation of Sphagnum steerei Andrus, 137

ANDRUS, R. E. (review). N. H. Holmgren (ed.), Winona H. Welch Festschrift. Brittonia 40: 113–199, 382

ANDRUS, R. E. Two new taxa of Sphagnum in section Cuspidata, 364

Aneura pinguis, 102

Anisomeridium biforme, 309+; nyssaegenum, new to Canada, 309+; willeyanum, 309+

Anomobryum conicum, 336; filiforme, 91, 336

Anomodon attenuatus, 98; minor, 98; rostratus, 98; rugelii, 99; viticulosus, 99

Antarctica, bryophyte oases in ablation valleys on Alexander Island, 45*

Anthoceros sp. 102

Antitrichia curtipendula, 358

Anzia colpodes, 309+

Aongstroemia jamaicensis, 335

Aphanorrhegma serrulatum, 99

Aptychella americana, *116*; colombica, *116*+, 119*; proligera, 116+, 338

Archidium acauloides, 214+; alternifolium, 214+; donnellii, 214+; tenerrimum, 214+

Arctic, species of Rinodina from, 106

Arctoa fulvella, 358

Arctoparmelia centrifuga, 309+

Arthonia leucopellaea, new to North America, 309+; phaeobaea, 171+, 188, new to eastern Canada, 309+; sp., 171, 188; spadicea, new to Canada, 309+

Arthopyrenia halodytes, *171*, *209*+

Arthothelium anastomosans, *309*+; dispersum, *309*+; ruanum, 309+

Arthrocormus schimperi, 28

Arthrorhaphis citrinella, 309+

Aschisma aethiopicum, 214+

Aspicilia cf. caesiocinerea, 309+; cinerea, 309+; laevata, 309+; sp., 171, 188; verrucigera, 309+

Asterella tenella, 102

Astomum ludovicianum, 99; muhlenbergianum, 214+; phascoides, 99

atmospheric pollution, lichen evidence for changes in Shenandoah National Park, Virginia, 21

Atrichum angustatum, 99; crispum, 99; ocrstedianum, 339; undulatum, 26, 99; undulatum var. altecristatum, 99; undulatum var. attenuatum, 99; undulatum var. oerstedianum, 99

Aulacomnium androgynum, 358; heterostichum, 99; palustre, 99

Aulacopilum glaucum, 191; hodgkinsoniae, 191; **schaeferi**, sp. nov., 191+, 192*; tumidulum, 191+

[1] Compiled by L. L. Sigal, assisted by R. O. Wadlington, Environmental Sciences Division, Oak Ridge National Laboratory, P.O. Box 2008, Oak Ridge, TN 37831-6038. Operated by Martin Marietta Energy Systems, Inc., for the U.S. Department of Energy under contract no. DE-AC05-84OR21400.

[2] New scientific names and new combinations are in boldface; illustrations are indicated by an asterisk, taxa discussed or mentioned several times in a paper are indicated by a plus sign; italicized page numbers indicate taxa in synonymy.

0007-2745/88/385–396$1.35/0

Bacidia atrogrisea, *309+*; chlorantha, 309+; cf. inundata, 309+; laurocerasi, 309+; naegelii, new to Canada, 309+; rubella, 309+; schweinitzii, 309+; scopulicola, 171+, 188; sp., 171, 188, 310+
Bactrospora dryina, 310+
Baeomyces fungoides, 310+; roseus, *310+*; rufus, 310+; sp., 310+
Barbella enervis, 32
Barbula arcuata, 30+; cancellata, 99; comosa, *37*; consanguinea, *37*; convoluta, 99; fallax, 99; icmadophila, 335; indica, 30+; javanica, 30+; louisiadum, 30; orientalis, *37*; unguiculata, 99
Bartramia angustifolia, 91, 116+, 336; defoliata, *116+*; longifolia, 91, 130; pomiformis, 99; thelioides, 91
Bazzania ambigua, 326+, mapped, 328; denudata, 102, 326+, mapped, 328; pearsonii, 326+, mapped, 328; pearsonii fo. **krajinai**, fo. nov., 326+, 331*; pearsonii fo. **schofieldii**, fo. nov., 326+, 331*; tricrenata, 102, 326+, mapped, 328; tricrenata var. **fulfordiae**, var. nov., 326+, 332*; tricrenata fo. **vittii**, fo. nov., 326+, 332*; trilobata, 102, 326+, mapped, 328
Belland, R. J. & W. B. Schofield. Pseudoleskea stenophylla Ren. & Card. *ex* Roell in eastern North America, 357
Biatorella moriformis, *310+*; ochrophora, *310+*; resinae, *310+*
Blasia pusilla, 102
Blepharostoma trichophyllum, 102
Botrydina vulgaris, *310+*
Bowers, F. D. *See* Timme, S. L. & F. D. Bowers, 368
Brachelyma subulatum, 99
Brachymenium acuminatum, 30; consimile, 131; daguense, 336; exile, 31; longicolle, 31; nepalense, 31
Brachythecium acuminatum, 99; austro-salebrosum, new to Peru, 91; curtum, 99; flexiventrosum, 338; oxycladon, 99; plumosum, 99, 338; rivulare, 99; rutabulum, growth rate of, 219+, 338; rutabulum var. flavescens, 99; rutabulum var. rutabulum, 99; salebrosum, 99; turgidum, 217; velutinum, 99
Braunfelsia dicranoides, 27; enervis, 27; longipes, 27; plicata, 27
Braunia secunda, 91
Brazil, a new species of Aulacopilum from, 191
Breen, Ruth Schornherst, a tribute to, 233
Breutelia arundinifolia, 31; brittoniae, 336; chrysea, 336; inclinata, 131, 336; karsteniana, 131; rhythidioides, 131; squarrosa, 131; subarcuata, 336; subdisticha, new to Colombia, 131; tomentosa, 336
Britain, growth rates of five bryophytes, 219
Brodo, I. M. *See* Gowan, S. P. & I. M. Brodo, 255
Brothera leana, 99
Bruchia flexuosa, 99
Bruggeman-Nannenga, M. A. *See* Pursell, R. A., M. A. Bruggeman-Nannenga & B. H. Allen, 202
Bryhnia graminicolor, 99; novae-angliae, 99
Bryoandersonia illecebra, 99
Bryoerythrophyllum jamesonii, 335; cf. recurvirostre, 47+; recurvirostrum, 99
Bryohaplocladium microphyllum, 99, 338, new to Colombia, 131; virginianum, 99
Bryohumbertia filifolia, 335; walkeri, 27
Bryoria bicolor, 310+; capillaris, 310+; friabilis, 310+; furcellata, 310+; fuscescens, 310+; implexa, 310+; nadvornikiana, 310+; salizinica, 310+; trichodes, 310+; trichodes subsp. americana, 310+; trichodes subsp. trichodes, 310+
Bryoxiphium norvegicum, 99
Bryum algovicum, 76+; alpinum, 75+; ambiguum, 31; amblyodon, 76+; andicola, 73+, 336; apiculatum, 31; archangelicum, 76+; arcticum, 76+; argenteum, 31, 74+, 99, 336; argenteum var. australe, 31; aubertii, *38*; bicolor, 99; blindii, 75+; bornholmense, 75+; caespiticium, *77+*, 99; calobryoides, 76+; calophyllum, 76+; canariense, 75+; capillare, 73+, 99; cognatum, 31; coronatum, 31, 336; crassirameum, *79*; cryophilum, 75+; cyclophyllum, 74+; dichotomum, 75+, 336; elegans, new to North America, 73+; erythroloma, 75+; flaccidum, 74+; gedeanum, 31; gemmascens, 76+; gemmilucens, new to North America, 73+; gemmiparum, 75+; graeffeanum, *38*; incrassatolimbatum, 73+; intermedium, 77+; junghuhnianum, 31; knowltonii, 76+; lanatum, 74+; lisae, 77+; lisae var. cuspidatum, 99; longisetum, 73+; maderense, 31; marratii, 76+; meesioides, 76+; microerythrocarpum, *79*; miniatum, 77+; muehlenbeckii, 75+; neilgherrense, 31; nitens, 31; oblongum, 77+; pallens, 74+; pallescens, 77+; plumosum, 31; porphyroneuron, 31; provinciale, 80; pseudotriquetrum, 47+, 74+, 99; purpurascens, 73+; pyriferum, 73+; radiculosum, new to North America, 73+; ramosum, excluded from North America, 73+; rubens, 75+; schleicheri, 76+; species in Western North America, 73; stirtonii, 77+; subapiculatum, 75+; subneodemense, 77+; tenuisetum, 75+; torquescens, 75+; turbinatum, 76+; uliginosum, 76+; violaceum, 75+; weigelii, 75+; wightii, 31; wrightii, 76+
Buck, W. R. Donnellia (Sematophyllaceae) resurrected and refound in Florida after 110 years, 134
Buck, W. R. *See* Anderson, L. E. & W. R. Buck, 113
Buck, W. R. *See* Delgadillo M., C. & W. R. Buck, 53
Buckley, A. & T. O. Hendrickson. The distribution of Cladonia perforata Evans on the southern Lake Wales Ridge in Highlands County, Florida, 354
Buellia aethalea, 310+; disciformis, 310+; parasema, *310+*; punctata, 310+; stillingiana, 310+; verruculosa, *310+*

Caliciales, key to species of northern Maine, 4
Calicium abietinum, 5+, 310+; glaucellum, 6+; hyperelloides, new to the United States, 367; parvum, 6+, 310+; salicinum, 6+, 310+; subquercinum, 7+; trabinellum, 7+, 310+; viride, 310+
Callicladium haldanianum, 99, 358
Callicosta bipinnata, 337; ramosissima, 337
Callicostella beccariana, 33; papillata, 33; prabaktiana, 33
Calliergonella cuspidata, 99
Caloplaca arenaria, 310+; aurantiaca, 188, *310+*; cerina, 310+; citrina, 171+, 188, 310+; discolor, 310+; ferruginea, 310+; flavorubescens, 310+; flavovirescens, 310+; granulosa, *171*; helicopsis, 178; holocarpa, 310+; cf. irrubescens, 311+; lactea, 188; marina, 178, 188; microthallina, 311+; murorum, 188; rosei, 188; spp., 171, 188; verruculifera, 171+, 188; cf. vittellinula, 311+
Calymperaceae, key to distinguish from Pottiaceae, 18
Calymperes, 18+; aeruginosum, 29; afzelii, 29+; angustatum, *38*; anisodictyon, *38*; beccarii, 29; bescherellei, *38*; constrictum, *29, 38*; crassinerve, 29+; cristatum, *38*; delessertii, *38*; **dixoconstrictum**, nom. nov., 29+; donnellii, *38*; dozyanum, 29+; erosum, 29+, 335; eutrichostomum, *38*; fasciculatum, 29+; fordii, *38*; hampei, *38*; heterophyllum, *38*; hyophilaceum, *38*; inaequifolium, *38*; kurzianum, *38*; lonchophyllum, 29+; longifolium, *38*; malayanum, *38*; moluccense, *38*; motleyi, 29; nicobarense, *38*; nietneri, *38*; nitidiusculum, 38; nymanii, *38*; orientale, *38*; orientale var. polytrichoides, *38*; palisotii, 29+; porrectum, 29+; punctulatum, 29; recurvifolium, *38*; repens, *38*; salakense, *38*;

schmidtii, 29+; serratum, 19, 29; setifolium, *38*; stenophyllum, 29; strictifolium, 29; subfasciculatum, *38*; subintegrum, *38*; taitense, 29+; tenerum, 29; vriesii, *38*; vriesii var. robustum, *38*
Calymperidium croceum, *38*; muelleri, *38*
Calymperopsis tjibodensis, 40
Calypogeja fissa, 102; muelleriana, 102; neesiana, 102; sullivantii, 102; trichomanis, 102
Calyptrochaeta remotifolia, 33+
Campylium chrysophyllum, 99; hispidulum, 99; polygamum, 47+, 99
Campylodontium flavescens, *38*
Campylopodium medium, 27+; euphorocladum, *38*; griffithii, *38*
Campylopus abbreviatus, *38*; argyrocaulon, 335; benedictii, new to Peru, 91; calodictyon, *38*; caudatus, *38*; comosus, 27+; crispifolius, 27; edithae, 335; ericoides, 27; exasperatus, 27+; exasperatus var. lorentzii, 27+; hemitrichus, 27; involutus, 27; lamellinervis, 335; laxitextus, 28+; macgregorii, 28; nivalis, 131, 335; pilifer, 99; pittieri, 335; richardii, 38, 335; ridleyi, *38*+; rosulatus, 335; serratus, 28; singapurensis, *38*; subconcolor, 335; tallulensis, 99; umbellatus, 28+; weberbaueri, 131
Campylostelium saxicola, 99
Canada, New Brunswick, lichens of Fundy National Park, 255
Candelaria concolor, 226
Candelariella aurella, 171, 187+, 311+; efflorescens, 311+; vitellina, 47+, 187, 311+; xanthostigma, 311+
CÁRDENAS S., A. Los musgos cleistocárpicos del Valle de México, México, 214
Catagonium brevicaudatum, 91, 337
Catillaria chalybeia, 171, 188, new to eastern Canada, 311+; globulosa, new to eastern Canada, 311+; sp., 171+, 188
Catinaria grossa, *311+*
Cephalozia bicuspidata subsp. bicuspidata, 102; bicuspidata subsp. lammersiana, 102; catenulata, 102; connivens, 102; lunulifolia, 102
Cephaloziella divaricata, 102; exiliflora, 47+; hampeana, 102
Ceratodon purpureus, 49, 99; purpureus fo. dimorphus, 235; stenocarpus, 131, 335
Cetraria aurescens, 311+; ciliaris, *311+*; halei, *311+*; hepatizon, 311+; islandica subsp. crispiformis, 311+; oakesiana, 311+; orbata, *311+*; pinastri, 311+; sepincola, 311+
Cetrariastrum catawbiense, new to Canada, 311+
Cetrelia chicitae, 311+; olivetorum, 311+
Chaenotheca brevipes, new to North America, 9+; brunneola, 7+, 311+; carthusiae, 7+; chrysocephala, 8+, 311+; consociata, 10+; debilis, 10+; ferruginea, 8+; furfuracea, 8+, 311+; hispidula, 9+; laevigata, 9+; lignicola, 10+; pusilla, 10+; rubescens, 11+; savonica, 11+; stemonea, 311+; trichialis, 9+; viridireagens, 11+
Chaenothecopsis lignicola, 311+; pusilla, 311+; subpusilla, *4*, *311+*
Chaetomitriopsis glaucocarpa, 33
Chaetomitrium borneense, 33, 39; ciliatum, 33; elongatum, 33; lanceolatum, 33; leptopoma, 34; muricatum, 34; nematosum, 34, 38; orthorrhynchum, 34; papillifolium, 34; perakense, 34; serratum, *38*; setosum, 34; torquescens, 34
Chiloscyphus pallescens, 102; polyanthos, 102
Chionoloma latifolium, 30
chlorophyll *b*, enhanced in Tomenthypnum nitens from simulated acid rain, 121
Chorisodontium mittenii, 335
chromosome numbers, in Riccia, 344
Chrysoblastella chilensis, 131
Chryso-hypnum diminutivum, 55
Chrysothrix candelaris, 171+, 188, new to Australia and South America, 360+; chlorina, 360; chrysophthalma, 360+; **granulosa**, sp. nov., 360+; pavonii, 360+; **placodioides**, sp. nov., 360+
CHURCHILL, S. P. Bryologia Novo Granatensis. Studies on the moss flora of Colombia I. New combinations, synonyms, and comments, 116
CHURCHILL, S. P. Bryologia Novo Granatensis. Studies on the moss flora of Colombia II. Further additions to Colombia and the Department of Antioquia, 130
CHURCHILL, S. P. & N. HOLLAENDER. Bryologia Novo Granatensis. Estudios de los flora de musgos de Colombia III. Catálogo de musgos del Departamento del Valle, 334
Cladina arbuscula, 311+; evansii, 354; mitis, 311+; rangiferina, 311+; stellaris, 311+; stygia, 311+; subtenuis, 354
Cladonia acuminata, 311+; anomaea, *311+*; bacillaris, 311+; boryi, 312+; botrytes, 312+; brevis, 312+; caespiticia, 312+; cariosa, 312+; carneola, 312+; cenotea, 312+; cervicornis, 312+; chlorophaea, 312+; coniocraea, 312+; conista, *312+*; cornuta subsp. cornuta, 312+; cornuta subsp. groenlandica, 312+; crispata, 312+; cristatella, 312+; cryptochlorophaea, 312+; deformis, 312+; digitata, 312+; farinacea, 312+; fimbriata, 312; floerkeana, 312+; furcata, 312+; gracilis, 312+; gracilis subsp. turbinata, 312+; grayi, 312+; humilis, 312+; leporina, 354; macilenta, 312+; macrophylla, 312+; maxima, 312+; merochlorophaea, 312+; metacorallifera, 47+, new to eastern Canada, 312+; multiformis, 312+; ochrochlora, 312+; pachycladodes, 354; perforata, distribution in Highlands County, Florida, 354, 355*; pityrea, *312+*; pleurota, 312+; prostrata, 354; pyxidata, 312+; ramulosa, 312+; rei, 313+; scabriuscula, 313+; squamosa var. squamosa, 313+; squamosa var. subsquamosa, 313+; subsetacea, 354; subulata, 313+; sulphurina, 313+; turgida, 313+; uncialis, 313+; verticillata, *313+*
Cladopodanthus pilifer, 28; speciosus, 38+
Claopodium nervosum, *38*; prionophyllum, 34
Clasmatodon parvulus, 99
Clastobryella asperrima, *38*; cuculligera, *38*; epiphylla, *38*; merrillii, *38*; ruficaulis, *38*
Clastobryophilum bogoricum, 35+; bogoricum var. laevibogoricum, *38*; rufoviride, 35+
Clastobryum caudatum, 35; cuculligerum, 35+; ruficaule, 35+; scalare, 35+
Clauzadea monticola, new to eastern Canada, 313+
Climacium americanum, 99; dendroides, 99
Cliostomum griffithii, 313+; sp., 313+
Coccocarpia cronia, *313+*; palmicola, 313+
Coelocaulon aculeatum, 313+
Collema cf. cristatum, 313+; fecundum, 171+, 187+; furfuraceum, 313+; subflaccidum, 313+; undulatum, 313+
Cololejeunea biddlecomiae, 102
Colombia, moss flora of Departamento del Valle, 334; studies on the moss flora of, 116, 130
Conardia compacta, 99
Coniocybe furfuracea, *4*, *313+*; gracilenta, *4*
Coniocybopsis arenarius, *4*
Conocephalum conicum, 102
Conotrema urceolatum, 313+
CRANDALL-STOTLER, B. *See* STOTLER, R. E. & B. CRANDALL-STOTLER, 56, 138, 236

Cribrodontium werneri, *38*
CROSBY, M. R. Recent literature on mosses—139, 151
Crossomitrium oerstedianum, 337; patrisiae, 337; wallisii, 337
CRUM, H. A new species of Aulacopilum from Brazil, 191
CRUM, H. (review). R. R. Ireland & G. Bellolio-Trucco, Illustrated Guide to Some Hornworts, Liverworts and Mosses of Eastern Canada, 250
CRUM H. (review). E. Nyholm, Illustrated Flora of Nordic Mosses. Fasc. 1. Fissidentaceae–Seligeriaceae, 250
Cryphaea glomerata, 99
Cryptogonium phyllogonioides, 33+
Ctenidiadelphus plumularia, 36+
Ctenidium falcifolium, *38*; lychnites, 36; malacodes, 339; molluscum, 99
CULBERSON, W. L. (review). W. J. Noble, T. Ahti, G. F. Otto & I. M. Brodo, A Second Checklist and Bibliography of the Lichens and Allied Fungi of British Columbia, 253
Cupressina malaccana, *38*
Cyathophorella burkillii, 33; tenera, 33
Cybebe gracilenta, 12+
Cyclodictyon blumeanum, 34; roridum, 337; rubrisetum, 337; subtortifolium, new to Colombia, 131
Cynodontium strumulosum, 357
Cyphelium lucidum, 12+

Daltonia aristofolia, 33; contorta, 33
DELGADILLO M., C. *See* SNIDER, J. A. & C. DELGADILLO M., 181
DELGADILLO M., C. & W. R. BUCK. Levierella (Fabroniaceae), a moss genus new to the Americas, 53
Dendriscocaulon umhausense, 313+
Dendrobazzania griffithiana, 326+
Dermatocarpon fluviatile, *313+*; luridum, 313+; miniatum, 313+; moulinsii, 313+; weberi, *313+*
Desmatodon obtusifolius, 99; porteri, 99
Desmotheca apiculata, 31
DEWEY, R. M. Electrophoretic studies in Riccia subgenus Riccia (Hepaticopsida: Ricciaceae), 344
Dichelyma capillaceum, 99
Dichodontium pellucidum, 99
Dicranella angustifolia, 335; brasiliensis, *38*; cerviculata, 99; coarctata, 28; heteromalla, 28, 99; cf. hilariana, 335; palustris, 358; perrottetii, 335; rufescens, 99; sp., 47+; vaginata, 335; varia, 99
Dicranodontium asperulum, 99; denudatum, 99; fleischerianum, 28+; nitidum, *38*; uncinatum, 28
Dicranoloma assimile, 28+; billardieri, 28+; blumii, 28; braunii, *38*; brevicapsulare, *38*; brevisetum, 28+; dives, 28; leucophyllum, *38*; perintegrum, 28; reflexifolium, *38*; reflexum, 28+; sumatranum, *38*
Dicranoweisia grimmiacea, 49
Dicranum condensatum, 99; flagellare, 99; frigidum, 335; fulvum, 93, 99; fuscescens, 99; **gregoryi**, sp. nov., 91+, 92*; majus, 358; megalophyllum, *38*; montanum, 99; polysetum, 99; scoparium, 99; spurium, 99; viride, 93, 99
Didymodon australasiae var. umbrosus, 18; cf. gelidus, 47+; sphagnoides, *38*
Dimerella diluta, *313+*; lutea, 313+; pineti 313+
Diphyscium cumberlandianum, 99; foliosum, 99; involutum 27; malayense, 27; mucronifolium, 27; rupestre, 27
Diplophyllum andrewsii, 102; apiculatum, 102
Diploschistes scruposus, 313+
Distichium capillaceum, 47*+
Distichophyllum cuspidatum, 34; gracilicaule, 34; malayense, 34; mittenii, 34; nigricaule, 34; pterygophylloides, 38; schmidtii, 34; sinuosulum, 34; spathulatum, 34; tortile, 34; ulukaliense, 34; undulatum, 34
Ditrichum difficile, 27+; flexicaule, 217; flexifolium, *38*; lineare, 100; pallidum, 100; pusillum, 100; submersum, 335
Donnellia **commutata**, comb. nov., 134+; florida, *134*; **lagenifera**, comb. nov., 134; **lageniformis**, comb. nov., 134; **laxiuscula**, comb. nov., 134; **longidens**, comb. nov., 134; **usambarica**, comb. nov., 134
Drepanocladus exannulatus, 338; fluitans, 100; uncinatus, 49
Drummondia prorepens, 100
Dryptodon patens, 358
Dumortiera hirsuta var. hirsuta, 102; hirsuta var. nepalensis, 102
Duthiella wallichii, 32

ECKEL, P. M. Taxa Minora Delineata, I. Ceratodon purpureus fo. dimorphus, 235
Ectropotheciella decrescens, 36+; distichophylla, 36
Ectropothecium buitenzorgii, 36+; callichroides var. elongatum, *38*; chamissonis, 36; dealbatum, 36; dentatum, *38*; eleganti-pinnatum, 36+; falcatulum, *38*; falcifolium, 36+; ichnotocladum, 36; incubans, 36+; incubans fo. scaberula 36+; intorquatum, 36; malacense, *38*; monumentorum, 37; moritzii, 37; moritzii var. stereodontoides, 37; penangense, 37; perminutum, 37; scaberulum, *38*; serratum, 37; siamense, 37; singapurense, 37; singapurianum, *38*; sparsipilum, 37; striatulum, 37; verrucosum, *38*; zollingeri, 37+
EGAN, R. S. Recent literature on lichens—132, 63;—133, 145;—134, 241
electrophoresis, studies in Riccia, 344
Encalypta, 18+; patagonica, 47+; procera, 47+
Endotrichella elegans, *38*; formosa, *38*; plano-marginata, *38*; polythrix, *38*
Entodon bandongiae, 34; brevisetus, 100; cladorrhizans, 100; hampeanus, 113+, 114*; macropodus, 100, 113+; seductrix, 100; sullivantii, 115; werneri, 34+
Entosthodon bonplandii, 336; jamesonii, 336; lindigii, 131
Ephebe lanata, 313+
Ephemeropsis tjibodensis, 33
Erioderma mollissimum, 313+
Eriopus remotifolius, *38*
Erpodium biseriatum, 31
Erythrodontium longisetum, 338
Eucamptodon wrayi, *38*
Eucladium verticillatum, 100
Eurhynchium hians, 100; pulchellum, 100; riparioides, 100
Eustichia longirostre, 336
Evernia mesomorpha, 314+
Everniastrum, *314+*
Exodictyon blumei, 28+; sullivantii, 28+
Exostratum blumei, *38*; sullivantii, *38*

Fabronia ciliaris, 338; ravenelii, 100
fen communities, effects of simulated acid rain on, 121
Fissidens adiantoides, 100; amblyotis, *39*; areolatus, 27; asplenioides, 39, 335; beccarii, 202+, 204*, 205*; bogoriensis, 27; brevilingulatus, 27; bryoides, 27, 100; bushii, 100; ceylonensis, 27+; crassinervis, 27; crenulatus var. elmeri, 27+; cristatus, 100; dissitifolius, new to Colombia, 334+; elegans, 39+; elmeri, *39*; exilis, 100; firmus, 27; fontanus, 100; geminiflorus, 27+; geminiflorus var. nagasakinus, *39*; grandifrons, 100; hollianus, 27+; hyalinus, 27, 100; hydropogon, 202+, 207*, 208*; intromarginatulus, *39*; javanicus, 27+; javanicus var. integrifolius, *39*; laxus, 27+; laxus var. **javensis**, comb. nov., 27; leptopelma, 27+; mangar-

evensis, 27; microcladus, 39; mitteni, *39*; mollis, 335; nobilis, 27; obtusifolius, 100; osmundoides, 100; papillosus, 27; polypodioides, 335; ravenelii, 100; repandus, 335; reticulosus, new to Colombia, 335; robinsonii, 27; siamensis, *39*; splachnobryoides, 27; stissotheca, 202+, 209*; stissotheca var. **brevicostatus**, var. nov. 202+, 212*; stissotheca var. **immersus**, stat. nov., 202+, 211*; stissotheca var. stissotheca, 202+; subangustus, *39*; subbasilaris, 100; subdiscolor, 27; subgenus Sarawakia, a taxonomic revision, 202; sylvaticus, 39; taxifolius, 100; teysmannianus, 27; weirii var. hemicraspedophyllus, 335; wichurae, 27; zippelianus, 27+; zollingeri, 27, 335
Fistulariella, 314+
Flavoparmelia baltimorensis, 21+; caperata, 226, 314+; sulfur and lead content in, 21+
Floribundaria floribunda, 32; pseudo-floribunda, 32; thuidioides, 32
Florida, distribution of Cladonia perforata in Highlands County, 354; Donnellia refound in, 134
Fontinalis dalecarlica, 100; filiformis, 100; flaccida, 100; novae-angliae, 100; sphagnifolia, 100; sullivantii, 100
Foraminella, *314+*
Forsstroemia ohioense, 103; trichomitria, 100, 103, 104
Fossombronia akiensis, *194*; angulosa, 193; australi-nipponica, 194; australis, 193+; brasiliensis, 194; caespitiformis, 193+; caledonica, *194*; capensis, 193; carinata, 194; corbulaeformis, 194; crassifolia, *193+*; crispa, 193+; cristata, 193+; cristula, 102, 195; crozalsii, 193; cubana, *195*; densilamellata, 193+; dentata, 195; dumortieri, *195*; echinata, 193; fernandeziensis, *193+*; fleischeri, *196*; forsythii, *193+*; foveolata, 102, 193+; gigantea, 196+; grandis, 193; gregaria, 196; grossepapillata, 196; hamato-hirta, *193+*; herzogii, 196; himalayensis, 196+; hispidissima, 196; husnotii, 197; incurva, 193; incurva var. tenera, 197; indica, 193+; integerrima, 197; integrifolia, 197; intermedia, 197; intestinalis, 193; japonica, *197*; lamellata, 197+; latifolia, 197; leucoxantha, 193+; leutzelburgiana, 197; levieri, *197*; loitlesbergeri, 197; longiseta, 197; lophoclada, 197; macounii, 198; macrocalyx, *198*; mauritanica, 198; mexicana, *193+*; microspora, 193; mittenii, 198; montaguensis, 193; naumannii, *198*; nigricaulis, 198; occidento-africana, *193+*; papillata, 193+; perpusilla, *199*; peruviana, 199; porphyrorhiza, 199; ptychophylla, 199; pulvinata, 199; pusilla, 193+; pusilla var. maritima, 195+; reticulata, 193; revisionary notes on, 193; rosulata, 199; rupestris, 195; salina, 199; spinifolia, 193+; spores, 193; stephanii, 200; subsaccata, 200; tenuifolia, 193+; texana, 193+; tuberifera, *200*; tumida, 193; tylimanthoidea, 200; verrucosa, 200; wattsii, 193; wondraczekii, 193+; wrightii, 193+; zeyheri, 200
Frullania brittoniae, 102; eboracensis, 102; inflata, 102; plana, 102; riparia, 102; squarrosa, 102; tamarisci subsp. asagrayana, 102
Funaria americana, 104; flavicans, 100, 104; hygrometrica, 30, 336; hygrometrica var. calvescens, 100; hygrometrica var. hygrometrica, 100
Fuscidea recensa, 114+

Gammiella merrillii, 35+
Garckea comosa, 27+; phascoides, *39*
Garovaglia angustifolia, 32; angustifolia fo. bogoriensis, 32+; aristata, 32; densifolia, *39*; elegans, 32+; elegans fo. formosa, 32+; plicata, 32; plicata var. splendida, 32; plicata subsp. undulata, 32; polythrix, *39*; powellii, 32; powellii subsp. densifolia, 39; powellii subsp. muelleri, 32; powellii subsp. muelleri var. plano-marginata, 32
Geocalyx graveolens, 103
GILMAN, B. *See* SLACK, N. G., C. RESCHKE & B. GILMAN, 217
Glossadelphus bilobatus, 35+; glossoides, 35; similans, 35; zollingeri, *39*
GOWAN, S. P. & I. M. BRODO. The lichens of Fundy National Park, New Brunswick, Canada, 255
GOWARD, T. Hypogymnia oceanica, a new lichen (Ascomycotina) from the Pacific Northwest of North America, 229
Graphis scripta, 114+
Grimmia alpicola var. alpicola, 100; alpicola var. rivularis, 100; apocarpa var. apocarpa, 100; apocarpa var. nigrescens, 100; apocarpa var. stricta, 100; donniana, 49; incurva, 358; laevigata, 100; longirostris, 336; maritima, 173; pilifera, 100; torquata, 358
Groutiella apiculata, 336; goniorrhyncha, 31+
growth rates, of five bryophytes in Britain, 219
Gyalecta convarians, 86; jenensis, new to eastern Canada, 314
Gyalidea lecideopsis var. convarians, rediscovered in Alaska, 86+, mapped, 89
Gymnostomiella vernicosa, 30
Gymnostomum aeruginosum, 100; angustifolium, 100; recurvirostre, 30; recurvirostrum, 100

Haemotomma cismonicum, 314+; elatinum, 314+; ochrophaeum, 314+
Hageniella nematosa, 35+
HALE, M. E., JR. *See* LAWREY, J. D. & M. E. HALE, JR., 21
HALE, M. E. (review). V. Wirth, Die Flechten Baden-Württembergs, 380
HAMADA, N. Individual variation in the contents of lichen substances in Ramalina nervulosa and R. subcomplanata, 95
Haplohymenium triste, 100
Harpalejeunea ovata subsp. integra, 103
Harpanthus scutatus, 103
Hedwigia ciliata, 100
Hedwigidium integrifolium, 336
Helicodontium capillare, 338
Helodium paludosum, 100
HENDRICKSON, T. O. *See* BUCKLEY, A. & T. O. HENDRICKSON, 354
Herbertus aduncus subsp. tenuis, 103
Herzogiella striatella, 100, 358; turfacea, 100
Heterocladium dimorphum, 358; macounii, 100, 357
Heterodermia obscurata, 314+; speciosa, 314+; squamulosa, new to Canada, 314+
Heterophyllium affine, 100, 338
Himantocladium arbuscula, *39*; exsertum, *39*; loriforme, 33; plumula, 33+; rugulosum, 33
HOLLAENDER, N. *See* CHURCHILL, S. P. & N. HOLLAENDER, 334
Holomitrium arboreum, new to Colombia, 131, 335
Homalia trichomanoides, 358
Homaliodendron exiguum, 33; flabellatum, 33; glossophyllum, *39*; intermedium, 33; javanicum, 33; microdendron, 33+; pinnatelloides, 33; scalpellifolium, 33; squarrulosum, 33
Homalotheciella subcapillata, 100
Homomallium adnatum, 100; incurvatum, 100
homosekikaic acid, in Ramalina nervulosa and R. subcomplanata, 95
HONG, W. S. The family Lepidoziaceae in North America west of the hundredth meridian, 326

Hookeria acutifolia, 100
Hookeriopsis cruegeriana, 337; diffusa, 337; purpureophylla, new to Colombia, 131; subfalcata, 337; undata, 131, 337
Horikawaea phyllogonioides, *39*
Huilia nigrocruenta, *314+*; panaeola, *314+*
Hydrogonium arcuatum, *39*; consanguineum, *39*
Hydrothyria venosa, new to eastern Canada, 314+
Hygrohypnum bestii, 357; closteri, 100; eugyrium, 100; micans, 100; ochraceum, 100
Hylocomium brevirostre, 100; splendens, 100
Hymenelia ceracea, 314+; lacustris, 314+
Hymenostomum edentulum, *39*; malayense, *39*
Hyophila commutata, *39*; dozy-molkenboeri, *39*; involuta, 30+, 93, 100, 335; javanica, 30; micholitzii, *39*
Hypnella diversifolia, 337
Hypnodendron arborescens, *39*; dendroides, 31+; diversifolium, 31+; junghuhnii, 31; reinwardtii, 31; subspininervium subsp. arborescens, 31+; wrayi, *39*
Hypnum amabile, 93; bogoricum, *39*; cupressiforme, 339; curvifolium, 100; fertile, 100; imponens, 100; lindbergii, 100; malaccanum, *39*; pallescens, 100, 358; polypterum, 339; pratense, 100, 104; sp., 471; subulatum, *39*; tavoyense, *39*
Hypocenomyce friesii, 314+; scalaris, 314+
Hypodontium, reclassified from Calymperaceae to Pottiaceae, 18+
Hypogymnia austerodes, 231; bitteri, 231; duplicata, 232; enteromorpha, 230+; heterophylla, 232; imshaugii, 232+; inactiva, 232; krogiae, 229+, 314+; metaphysodes, 231; mollis, 231; occidentalis, 230+; **oceanica**, sp. nov., 229+; physodes, 231, 314+; pseudophysodes, 229+; pulverata, 232; rugosa, 231; subobscura, 231; tubulosa, 231, 314+; vittata, 231, 314+
Hypopterygium aristatum, 34; javanicum, *39*; rigidulum, *116+*, 119*; sylvaticum, *116+*, 119*; tamariscinum, 116+, 338; tenellum, 34
Hypotrachyna revoluta, new to eastern Canada, 314+

Icmadophila ericetorum, 315+
Imshaugia aleurites, 315+
Index of Atmospheric Purity (IAP), 223
Ionaspis odora, new to eastern Canada, 315+; sp., 171+, 188
Isodrepanium lentulum, 337
Isopterygiopsis muelleriana, 100
Isopterygium albescens, 37; arquifolium, 37; bancanum, 37; chrismarii, 339; constrictum, *39*; distichaceum, 100; elegans, 100, 104, 358; laxissimum, 37; minutirameum, 37; minutirameum var. tonkinense, 37; pohliaecarpum, 37+; subalbescens, 37; tenerum, 93, 100; textorii, *39*

Jamesoniella autumnalis, 103
Jubula pennsylvanica, 103
Julella fallaciosa, 315+
Jungermannia atrovirens, 103; crenuliformis, 103; gracillima, 103; hyalina, 103; pumila, 103

Kentucky, a list of bryophytes of, 98
Kindbergia praelonga, 93
Kingiobryum paramicola, 18
Kurzia pauciflora, 326+, mapped, 328; sylvatica, 103, 326+, mapped, 328; trichoclados, 326+, mapped, 328

Lasallia papulosa, 315+
LAWREY, J. D. (review). D. H. S. Richardson (ed.). Biological Indicators of Pollution, 70
LAWREY, J. D. & M. E. HALE, JR. Lichen evidence for changes in atmospheric pollution in Shenandoah National Park, Virginia, 21
Lead, accumulation in Flavoparmelia baltimorensis, 21+
Lecanactis abietina, 315+; dryina, *315+*; dryophila, *315+*
Lecania aipospila, 188; spp., 171+, 188
Lecanora actophila, new to North America, 315+; albescens, 171+, 188, 315+; bockii, *315+*; caesiorubella subsp. caesiorubella, 315+; chlarona, *315+*; cinereofusca var. cinereofusca, 315+; contractula, 315+; dispersa, 171+, 188+, 315+; expallens, 315+; fuliginosa, 315+; fuscidula, new to Canada, 315+; glabrata, 315+; grantii, 171+, 188, 315+; hageni, 188, 315+; helicopsis, 188; hybocarpa, 315+; intricata, 315+; mutabilis, *315+*; orae-frigidae, 315+; poliophaea, 188; polytropa, 315+; polytropa var. intricata, *315+*; pulicaris, 315+; subrugosa, 315+; symmicta, 315+; symmicta var. sorediosa, *315+*; symmictera, *315+*; thysanophora, 315+; cf. thysanophora, 315+; umbrina, 315+
Lecidea aeruginosa, *315+*; albofuscescens, 315+; atrofulva, 315+; auriculata, 315+; berengeriana, 315+; brunneofusca, 316+; cinnabarina, 316+; cyrtidia, 316+; elabens, 316+; epixanthoidiza, 316+; erratica, 316+; granulosa, *316+*; helvola, 316+; latypea, *171*; lucida, *316+*; monticola, *316+*; oligotropha, *316+*; paraclitica, *316+*; plana, 316+; plebeja, new to eastern Canada, 316+; speirea, 316+; spp., 171, 188; tornoënsis, *316+*; tumida, *316+*; uliginosa, *316+*; varians, *316+*; vernalis, 316+; viridescens, 316+
Lecidella scabra, 171, 188; stigmatea, 171+, 188, 316+; subincongrua, 171+, 188
Leiomela bartramioides, 336; lopezii, 336
Lejeunea cavifolia, 103; laetivirens, 103; lamacerina subsp. gemminata, 103; ruthii, 103; ulicina subsp. bullata, 103; ulicina subsp. ulicina, 103
Lemmholemma myriococcum, 316+
Lepidopilum aubertii, 337; brevipes, 337; erubescens, 337; excelsum, 337; frondosum, 337; perlaxum, 338; permarginatum, 338; radicale, 338; scabrisetum, 338; surinamense, 131; tortifolium, 338
Lepidozia filamentosa, 326+, mapped, 328; reptans, 103, 326+, mapped, 328; sandvicensis, 326+, mapped, 328
Lepidoziaceae, in North America, 326
Lepraria finkii, 316+; membranacea, 316+; neglecta, *316+*; cf. neglecta, 47+; sp., 171+, 188; zonata, 316+
Leptobryum pyriforme, 100
Leptodontium tricolor, 19; viticulosoides, 93, 336; viticulosoides var. exasperatum, new to Peru, 93
Leptogium corniculatum, 171, 188; cyanescens, 316+; laceroides, 316+; lichenoides, 316+; minutissimum, 171+, 188; palmatum, *171*; plicatile, 171, 188; rivale, 171, 188; saturninum, 316+; tenuissimum, 171, 188
Leptorhaphis epidermidis, 316+
Leptotheca boliviana, 336
Lepyrodon tomentosus, 337
Lescuraea saxicola, 357+; stenophylla, 357+
Leskea angustata, 131; gracilescens, 100; obscura, 100; polycarpa, 100
Leskeodon longipilus, 131; wallisii, 338
Leucobryum aduncum, 28; albicans, 335; albidum, 100; antillarum, 335; bowringii, 28; candidum, 28; chlorophyllosum, 28; crispum, 335; cyathifolium, 28; giganteum, 335; glaucum, 100; javense, 28+; martianum, 335; microleucophanoides, 28; neilgherrense, 28; pulchrum, 28+; sanctum, 28; scabrum, 28; scalare, 28; subsanctum, 28; stellatum, *39*; sumatranum, *39*
Leucodon brachypus, 104; brachypus var. andrewsianus, 101; brachypus var. brachypus, 101; julaceus, 101
Leucolejeunea clypeata, 103; unciloba, 103

Leucoloma malayanum, 39; molle, 28; pallidum, 39; serrulatum, 335; walkeri, 28
Leucomium aneurodictyon, *39*; strumosum, 37+, 338
Leucophanes aciculare, *39*; albescens, 28; candidum, 28+; candidum var. densifolium, 39+; densifolium, *39*; octoblepharoides, 29; pugioniforme, *39*
Levierella perserrata, new to Mexico and the Americas, 53*+
lichen communities, effects of O_3 and SO_2, 223
Lindbergia brachyptera, 101
Lobaria pulmonaria, 316+; quercizans, 316+; scrobiculata, 316+
Lopadium disciforme, 316+
Lophocolea bidentata, 103, growth rate of, 219+; cuspidata, 103; heterophylla, 103
Lophozia bicrenata, 103; incisa, 103
Lopidium struthiopteris, 34+; trichocladon, 34

Macrohymenium muelleri, 35; strictum, 35
Macromitrium blumei, 31; brevirete, 39; cirrosum, 131, 336; cumingii, 32; cuspidatum, 32; goniorrhynchum, *39*; incurvifolium, 32; longicaule, 32; longifolium, 336; micropoma, 32; minutum, 32; ochraceum, 32; orthostichum, 32; portoricense, 131; reinwardtii, 32; salakanum, 32; scoparium, 336; semipellucidum, 32; serrulatum, 336; stellatum, 131; subtile, 32; trachypodium, 336; zollingeri, 32
Macrothamnium macrocarpon, 37
MAGILL, R. E. (review). J.-P. Frahm & W. Frey, Moosflora (Second edition), 159
MAGILL, R. E. (review). B. M. Murray, Illustrated Moss Flora of Arctic North America and Greenland. 3. Andreaeobryaceae–Tetraphidaceae, 252
Maine, Caliciales in, 4
malate dehydrogenase (MDH), variability in Riccia, 344
Malaysia, peninsular, checklist for mosses of, 24, map, 26
Marchantia polymorpha, 103
MARINO, P. C. The North American distribution of the circumboreal species of Splachnum and Tetraplodon, 161
Marsupella emarginata, 103; sphacelata, 103
Mastopoma impolitum, 39; papillatum, 39; perundulata, 35
MAYRHOFER, H. & J. W. SHEARD. Four notable saxicolous species of the lichenized ascomycete genus Rinodina from the Arctic, 106
MCCUNE, B. Lichen communities along O_3 and SO_2 gradients in Indianapolis, 223
MEDLEY, M. *See* SNIDER, J. A., S. M. STUDLAR & M. MEDLEY, 98
Meesia lavardei, 336
Megalaria grossa, 317+
Meiotheciopsis, *134*+; commutata, *134*
Meiothecium, 134+; hamatum, 35; jagorii, 35; microcarpum, 35+; microcarpum var. lineolatum, 35; scaberulum, 39; tenerum, 134
Melanelia disjuncta, 317+; exasperatula, 317+; glabratula, 317+; halei, 317+; olivacea, 317+; panniformis, 317+; septentrionalis, 317+; sorediata, 317+; sorediosa, *317*+; subaurifera, 171, 188, 317+
Menegazzia terebrata, 317+
Mesonodon flavescens, 34+, 338
Meteoridium remotifolium, 337
Meteoriella soluta, 32
Meteoriopsis ancistrodes, *39*; reclinata, 32; reclinata var. ancistrodes, 33+; reclinata var. subreclinata, 32
Meteorium atro-lutea, *116*+, 119*; illecebrum, 116+; miquelianum, 33
4′-*O*-methylnorhomosekikaic acid, in Ramalina subcomplanata, 95
Metzgeria conjugata, 103; crassipilis, 103; furcata, 103; myriopoda, 103
Mexico, cleistocarpous mosses in the Valley of Mexico, 214
Mexico, Pleuridium aurantiacum, a new species from, 181
Micarea bacidiella, 317+; globulosella, 317+; hedlundii, new to North America, 317+; lutulata, new to North America, 317+; melaena, 317+; misella, 317+; peliocarpa, 317+; prasina, 317+; sylvicola, 317+; viridescens, *317*+
Microcalicium arenarium, 12+; disseminatum, 12+
Microcampylopus khasianus, 28; laevigatum, 28
Microdus brasiliensis, 28+; macromorphus, 28; miquelianus, 28
Microglaena corrosa, *317*+
Micromitrium synoicum, 101
Mielichhoferia macrocarpa, 357
MILLER, N. G. (review). C. Oostendorp, The Bryophytes of the Palaeozoic and the Mesozoic, 69
MILLER, N. G. (review). J. C. Ritchie, Postglacial Vegetation of Canada, 249
Mittenothamnium lehmannii, 339; loriforme, 116+; reptans, 339
Mitthyridium cardotii, 29; constrictum, 29; crassum, 29; fasciculatum, 29+; flavum, 29+; jungquilianum, 29; louisiadum, *39*; luteum, 29; manii, *39*; pungens, 29+; repens, 29+; repens var. **minor**, comb. nov., 29; subluteum, 29; undulatum, 29+; wallisii, 30+
Mniodendron curtisii, *39*; divaricatum, *39*; mittenii, *39*+
Mniomalia semilimbata, 31
Mnium affine, 101; affine var. ciliare, 101; affine var. rugicum, 101; cuspidatum, 101; hornum, 101, 104; integrum, *39*; longirostre, *39*; longirostrum, 101; lycopodioides, 101; marginatum, 101; medium, 101; punctatum var. appalachianum, 101; punctatum var. elatum, 101; punctatum var. punctatum, 101; rhynchophorum, *39*; spinulosum, 104; stellare, 101; succulentum, *39*; thomsonii, 101
MOHAMED, H. & W. D. REESE. Syrrhopodon mammillosus C. Muell., 51
MOHAMED, H. & B. C. TAN. A checklist of mosses of peninsular Malaysia and Singapore, 24
Mosigia cf. gibbosa, new to Canada, 317+
MURRAY, B. M. *See* THOMSON, J. W. & B. M. MURRAY, 86
Mycocalicium parietinum, *317*+; savonicum, *4*; subtile, 13+, 317+
Mycoporum compositum, 317+; pycnocarpum, *317*+
Mykoblastus affinis, new to eastern Canada, 317+; sanguinarius, 317+; tornoënsis, 317+
Myurella siberica, 101
Myurium pseudo-rufescens, *39*; rufescens, *39*; subnitens, *39*

Nadvornikia hawaiensis, new to the United States, 367
Nardia lescurii, 103
NASH, T. H., III (review). J. Middelborg & J. Mattson, Crustaceous lichenized species of the Caliciales in Norway. Sommerfeltia 5: 1–70, 381
Neckera pennata, 101, 358; spruceana, new to Colombia, 337
Neckeropsis andamana, 33; disticha, 337; exserta, 33+; fimbriata, 33; gracilenta, 33+; lepineana, 33; penicillata, *39*; submarginata, 33
Neolindbergia semitorta, 32
Nephroma arcticum, 317+; bellum, 317+; helveticum, 317+; parile, 317+; resupinatum, 317+

New Guinea, the moss genus Papillaria in, 341
New York, Scorpidium turgescens rediscovered, 217
Normandina pulchella, new to eastern Canada, 317+
North America, species of Hypogymnia from the Pacific Northwest, 229; the family Lepidoziaceae in, 326; western species of Bryum, 73
Notothylas orbicularis, 103
Nowellia curvifolia, 103

Ochrobryum **herzogianum**, nom. nov., 29+; microphyllum, *39*
Ochrolechia androgyna, 318+; pseudopallescens, 318+; rosella, 318+; cf. rosella, 318+
Octoblepharum albidum, 29, 335; cocuiense, 335
Odontoschisma denudatum, 103; prostratum, 103
Oedicladium **pseudo-rufescens**, comb. nov., 32+; rufescens, 32+; subnitens, 32+
Oedipodium griffithianum, 358
Oligotrichum hercynicum, 358
Omphalina ericetorum, *318+*
Opegrapha saxicola, 171+, 188
Orthodontium gracile, 93; infractum, 31; pellucens, 131
Orthomnion dilatatum, 31+
Orthomniopsis dilatata, *39*
Orthorrhynchium philippinense, *39*; ridleyi, *39*
Orthostichidium guyanense, 337
Orthostichopsis tetragona, 337
Orthotrichum anomalum, 101; gymnostomum, 358; ohioense, 101; pumilum, 101; pusillum, 101; sordidum, 101, 358; stellatum, 101; strangulatum, 101
Osculatia columbica, 336
Oxystegus tenuirostris, 336
ozone, effects on lichen communities in Indianapolis, 223

Pallavicinia lyellii, 103
Pannaria ahlneri, 318+; conoplea, 318+; leucophaea, 318+; pezizoides, 318+; pityrea, *318+*; rubiginosa, 318+
Papillaria crocea, 33, new to New Guinea, 341+, 342*, mapped, 342; flexicaulis, 341+, 342*, mapped, 342; fuscescens, 33+, 341+, mapped, 342; fuscescens var. rigidicaulis, *39*; intricata, 341; leuconeura, new to New Guinea, 341+, 342*; nigrescens, 337; penicillata, 337
Paraleucobryum longifolium, 101
Parmelia appalachensis, *318+*; arnoldii, *318+*; aurulenta, *318+*; caperata, *318+*; centrifuga, *318+*; conspersa, *318+*; crinita, *318+*; cumberlandia, *318+*; disjuncta, *318+*; exasperatula, *318+*; fertilis, 318+; galbina, *318+*; glabratula, *318+*; halei, *318+*; olivacea, *318+*; omphalodes, 178, 318; panniformis, *318+*; perlata, *318+*; revoluta, *318+*; rudecta, *318+*; saxatilis, 178, 188, *318+*; saxatilis var. divaricata, *318+*; septentrionalis, *318+*; sorediosa, *318+*; squarrosa, *318+*; subaurifera, *171, 318+*; subrudecta, *318+*; sulcata, 318+; taractica, 318+
Parmeliella corallinoides, *318+*; triptophylla, 318+
Parmelina aurulenta, 226, 318+; galbina, 318+
Parmeliopsis aleurites, *318+*; ambigua, 318+; hyperopta, 318+
Parmotrema arnoldii, 318+; chinense, 318+; crinitum, 318+; perlatum, *319+*
Pelekium bifarium, 34+; velatum, 34+
Pellia endiviifolia, 104; epiphylla, 103; megaspora, 104; neesiana, 103
Peltigera aphthosa, 319+; canina, 319+; degenii, 319+; didactyla, 319+; elisabethae, 319+; evansiana, 319+; horizontalis, 319+; leucophlebia, 319+; membranacea, 319+; neckeri, 319+; neopolydactyla, 319+; polydactyla, 319+; ponojensis, 319+; praetextata, 319+; rufescens, 319+; scabrosa, 319+; spuria, *319+*; venosa, 319+
peroxidase (PER), variability in Riccia, 344
Pertusaria alpina, 319+; amara, 319+; consocians, 319+; globularis, new to Canada, 319+; leucostoma, 319+; macounii, 319+; multipunctoides, 319+; neoscotica, 319+; ophthalmiza, 319+; propinqua, new to Canada, 319+; rubefacta, 319+; subobducens, 319+; trachythallina, 319+; velata, 319+; waghornei, 319+
Peru, contributions to the moss flora of, 86
Petalophyllum ralfsii, 194
Phaeocalicium compressulum, 13+; curtisii, 13+; polyporaeum, 14+; populneum, 14+
Phaeoceros laevis, 103
Phaeographis inusta, 319+
Phaeophyscia adiastola, 319+; ciliata, 319+; orbicularis, 171+, 188; pusilloides, 226, 319+; rubropulchra, 226, 319+; sciastra, 171+, 189, 319+
Philonotis angulata, 336; capillaris, 101; fontana, 101; fontana var. caespitosa, 101; glaucescens, 336; gracillima, 336; hastata, 31+; laxissima, *39*; longiseta, 101; marchica, 101; mollis, 31; muhlenbergii, 101; secunda, 31; speciosa, 31; turneriana, 31; uncinata, 336
Phlyctis agelaea, 320+; argena, 320+
Phyllodrepanium falcifolium, 336
Phyllogonium fulgens, 337; viscosum, 337
Physcia adscendens, 171, 189, 226, 320+; aipolia, 320+, caesia, 47+, 171+, 178+, 320+; dubia, 320+; millegrana, 226, 320+; orbicularis, *171*; phaea, 320+; sciastra, *171*; stellaris, 226, 320+; tenella, 171, 189, 320+
Physcomitrium immersum, 101; pyriforme, 30, 101
Physconia detersa, 226, 320+; muscigena, 47+
Phytoconis ericetorum, 320+
Pike, D. C. *See* Scott, G. A. M. & D. C. Pike, 193
Piloecium pseudo-rufescens, *39*
Pilopogon longirostratus, 335
Pilotrichella flexilis, 337; pentasticha, 337; perakensis, *39*
Pinnatella alopecuroides, 33; ambigua, 33; anacamptolepis, 33; kuehliana, 33; lingulata, 33; microptera, 33; mucronata, 33
Placynthiella icmalea, 320+; oligotropha, new to Canada, 320+; uliginosa, 320+
Placynthium flabellosum, 320+; nigrum, 320+; sp., 171, 189
Plagiochila austini, 103; porelloides, 103; sullivantii, 103
Plagiomnium integrum, 31+; rhynchophorum, 31, 336; rostratum, 31+; succulentum, 31+; undulatum, growth rate of, 219+
Plagiothecium, 47+; cavifolium, 101; denticulatum, 101, 338; laetum, 101; miquelii, *39*; neckeroideum, 34; simonovii, 49
Platismatia glauca, 320+; tuckermanii, 320+
Platydicta confervoides, 101; subtile, 101
Platygyrium repens, 101
Platyhypnidium aquaticum, 338
Pleuridium **aurantiacum**, sp. nov., 181+, 182*; **lindigianum**, comb. nov., 116+, 119*; sp., 214+; subulatum, 101, 214+; sullivantii var. mexicanum, 214+
Pleurochaete squarrosa, new to Mississippi, 368+
Pleurozium schreberi, 101
Pogonatum brachyphyllum 101; campylocarpum, 93; cirratum, 26+; convolutum var. cirratum, *39*; flexicaule, 26; junghuhnianum, 26; macrophyllum, 26; neesii, 26; neglectum, 339; oligodus, 339; pensilvanicum, 101; piliferum, *39*; semipellucidum, 339; spuriocirratum, 26; subtortile, 26; subtortile var. teysmannianum, 26; teysmannianum, *39*
Pohlia andalusica, 358; cruda, 47+; elongata, 41; lescu-

riana, 101; nutans, 49, 101; papillosa, 131; wahlenbergii, 101
Polyblastia **cucurbitula**, sp. nov., 86+, 87*, 88*, mapped, 89; cupularis, 90; fertilis, 90; hyperborea, 90; sp., 171+, 189; terrestris, 90+; theleodes, 90, new to eastern Canada, 320+
Polyblastiopsis fallaciosa, *320*+
Polychidium muscicola, new to eastern Canada, 320+
Polytrichadelphus aristatus, 339; ciliatus, 339; longisetus, 339; purpureus, 339
Polytrichum commune, 101; formosum, 101; juniperinum, 101, 339; ohioense, 101; pallidisetum, 101; piliferum, 101
Porella pinnata, 103; platyphylla, 103; platyphylloidea, 103
Porotrichodendron robustum, 337; superbum, 337
Porotrichum expansum, 338; korthalsianum, 338; longirostre, 338; mutabile, 338
Porpidia albocaerulescens, 320+; cinereoatra, 320+; crustulata, 320+; diversa, 320+; macrocarpa, 320+; soredizodes, 320+; speirea, 320+; tahawasiana, 320+; tuberculosa, 320+
Potamium recurvifolium, 338
Pottia truncata, 100
Pottiaceae, key to distinguish from Calymperaceae, 18
Preissia quadrata, 103
Prionodon densus, 337; fusco-lutescens, 337
Protoblastenia cinnabarina, *321*+; rupestris, 321+
Protothelenella cf. corrosa, new to eastern Canada, 321+
Pseudephebe minuscula, 47+
Pseudevernia cladonia, new to Canada, 321+
Pseudobarbella ancistroides, *39*; assimilis, *39*; attenuata, 33+
Pseudocyphellaria crocata, 321+
Pseudoleskea patens, 357+; radicosa, 357; stenophylla, 357+, mapped, 358
Pseudoparmelia caperata, *321*+
Pseudoscleropodium purum, growth rate of, 219+
Psilolechia lucida, new to eastern Canada, 321+
Pterobryon densum, 117+, 119*; excelsum, 116+, 119*; hornschuchii, *117*; lorentzii, *117*+; panamense, *117*; stolonaceum, *117*; subangustifolium, *117*
Pterobryopsis crassicaulis, 32; gedehensis, 32; stolonacea, 117
Ptilidium pulcherrimum, 103
Ptilium crista-castrensis, 101
Ptychomitrium drummondii, 101; incurvum, 101
Punctelia appalachensis, 321+; rudecta, 226, 321+; subrudecta, 226, 321+
PURSELL, R. A., M. A. BRUGGEMAN-NANNENGA & B. H. ALLEN. A taxonomic revision of Fissidens subgenus Sarawakia (Bryopsida: Fissidentaceae), 202
Pursellia phyllogonioides, *39*+
Pycnothelia papillaria, 321+
Pylaisiadelpha recurvans, 101; tenuirostris, 101
Pylaisiella intricata, 101; selwynii, 101
Pyrenocollema halodytes, 171+, 189, 321+; imshaugii, 189; spp., 171+, 189
Pyrenopsis pulvinata, 217
Pyrenula laevigata, 321+; neglecta, *321*+; pseudobufonia, 321+
Pyrrhobryum latifolium, 31+; longiflorum, *40*; spiniforme, 31+, 336; spiniforme var. badakense, *40*
Pyrrhospora varians, 321+
Pyxine caesiopruinosa, 226; sorediata, 226, 321+

Racelopus pilifer, 27+
Racomitrium aciculare, 101; crispipilum, 336; heterostichum var. microcarpon, 101; lanuginosum, 336
Racopilum cuspidigerum, 32+; orthocarpum, 32; schmidii, 32, *40*; spectabile, 32; tomentosum, 336
Radula complanata, 103; obconica, 103; sullivantii, 103; tenax, 103
Ramalina americana, 321+; dilacerata, 321+; farinacea, 171, 189, 321+; hypoprotocetrarica, *321*+; intermedia, 321+; minuscula, *321*+; nervulosa, lichen acids in, 95; pollinaria, 321+; roesleri, 321+; siliquosa, 95; subcomplanata, lichen substances in, 95; thrausta, 321+
RAVEN, P. H. (review). L. Margulis & K. V. Schwartz, Five Kingdoms: An Illustrated Guide to the Phyla of Life on Earth, Second Edition, 382
REESE, W. D. Editorial, 1
REESE, W. D. Rhizoid gemmae in Weissia controversa Hedw., 184
REESE, W. D. Ruth Schornherst Breen, 233
REESE, W. D. *See* MOHAMED, H. & W. D. REESE, 51
REESE, W. D. & R. H. ZANDER. Comparison of Calymperaceae with Pottiaceae, 18
REESE, W. D. & R. H. ZANDER. Typification of Serpotortella Dixon (Musci), 136
RESCHKE, C. *See* SLACK, N. G., C. RESCHKE & B. GILMAN, 217
Rhabdoweisia crispata, 101; fugax, 93
Rhacocarpus purpurascens, 336
Rhacopilopsis trinitensis, new to Colombia, 131
Rhaphidostegium caespitosum, *40*; complanatulum, 40; densirameum, 40; microcladum, *40*; saproxylophilum, *40*; subpinnatum, 40+; tristiculum, *40*
Rhaphidostichum boschii, *40*; bruchii, 35+; luxurians, 35+; malayanum, 35+; surculare, 35+
Rhizocarpon badioatrum, 321+; cinereovirens, 321+; cf. concentricum, 321+; disporum, 171+, 189; geographicum, 49, 321+; grande, 321+; hochstetteri, 321+; lavatum, 321+; lecanorinum, 322+; macrosporum, new to eastern Canada, 322+; obscuratum 322+; oederi, new to eastern Canada, 322+; petraeum, *322*+; plicatile, 322+; subgeminatum, new to Canada, 322+; submodestum, 322+; tetramerum, *322*+; tinei, *322*+; umbilicatum, new to eastern Canada, 322+
Rhizogonium badakense, *40*; graeffeanum, 31; latifolium, *40*; lindigii, 336; longiflorum, *40*; novae-hollandiae, 31; spiniforme, *40*; spiniforme var. badakense, *40*
rhizoid gemmae, in Weissia controversa, 184+
Rhizomnium magnifolium, 358
Rhizoplaca melanophthalma, 47+
Rhodobryum aubertii, 31+; beyrichianum, 336; giganteum, 31; graffeanum, *40*; grandifolium, 336; ontariense, 31; roseum, 101
Rhynchostegiopsis complanata, 131; costaricensis, 131; flexuosa, new to Colombia, 131; planifolia, 131; tunguraguana, 131
Rhynchostegium celebicum, 34; javanicum, 34; scariosum, 338; serrulatum, 101, new to Colombia, 338, 358
Rhytidiadelphus loreus, 358; novogranatensis, *116*+, 119*
Rhytidium rugosum, 101
Riccardia chamedryfolia, 103; latifrons, 103; multifida, 103; palmata, 103
Riccia albida, 345+; austinii, 346+; beyrichiana, 346+; bifurca, 346+; californica, 346+; campbelliana, 345+; dictyospora, 344+; donnellii, 346+; eldeeniae, 344+; fluitans, 103, 346+; hirta, 344+; mcallisteri, 344+; membranacea, 103; nigrella, 346+; sorocarpa, 103, 346+; sullivantii, 103; trichocarpa, 346+
Ricciocarpus natans, 103
Rigodium toxarion, 338
RINCON, E. Effect of initial shoot size on growth rates of five British grassland bryophytes, 219
Rinodina arnoldii, new from the Arctic, 106+, spores,

109*, 110*; ascociscana, 322+; bischoffii, 107; calcigena, 106+, mapped, 108; castanomela, 106+; castanomelodes, 106+; endophragmia, 106+, mapped, 108, spores, 109*; gennarii, 171+, 189; iodes var. iodes, 109+, iowensis, 322+; lecanorina, 109; magnusoni, 322+; milliaria, 322+; milvinia, 111+; occidentalis, 107+; ochracea, 106+ salina, *171*; straussii, 106+; subminuta, 322+
ROCHEFORT, L. & D. H. VITT. Effects of simulated acid rain on Tomenthypnum nitens and Scorpidium scorpioides in a rich fen, 121
RYAN, B. D. Marine and maritime lichens on serpentine rock on Fidalgo Island, Washington, 186
RYAN, B. D. Zonation of lichens on a rocky seashore on Fidalgo Island, Washington, 167

salazinic acid, in Ramalina subcomplanata, 95
Sarawakia, a subgenus of Fissidens, 202
Sarconeurum glaciale, 49
Sarea resinae, 322+
Sauloma microcarpa, *40*
Scapania nemorosa, 103; undulata, 103
Schistidium antarcticum, 47+; trichodon, 358
Schistomitrium apiculatum, 29; mucronifolium, 29; nieuwenhuisii, 29; sparei, 29
Schistostega pennata, 358
Schizomitrium **acutifolium**, comb. nov., 116+, 338; colombicum, 120, 338; pallidum, 338; **plicatulum**, comb. nov., 116, 338; rivulare, 120
Schizymenium campylocarpum, 93, 336
Schliephackea meteorioides, 335
Schlotheimia, new to Colombia, 131; angustata, 131+; rugifolia, 132+; torquata, 132+
SCHOFIELD, W. B. *See* BELLAND, R. J. & W. B. SCHOFIELD, 357
Schwetschkeopsis fabronia, 101
Sciaromium lescurii, 101
Sclerohypnum riparium, 34
Scoliosporum chlorococcum, 322+; umbrinum, 322+
Scopelophila cataractae, 101
Scorpidium scorpioides, 101, effects of simulated acid rain on, 121; turgescens, rediscovered in New York State, 217+, 218*
SCOTT, G. A. M. & D. C. PIKE. Revisionary notes on Fossombronia, 193
sekikaic acid, in Ramalina nervulosa and R. subcomplanata, 95
Seligeria brevifolia, 358; pusilla, 101; tristichoides, 358
SELVA, S. B. The Caliciales of northern Maine, 2
Sematophyllum adnatum, 101, 135+; bogoricum, *40*; braunii, *40*; caespitosum, 40+; cuspidiferum, 338; decipiens, *40*; demissum, 102; dicranolomoides, *40*; flavidum, 338; galipense, 338; hyalinum, *40*; lindigii, 338; longicuspis, *40*; marylandicum, 102; microcladiellum, 35+; obscurum, *40*; perpapillosum, *40*; rigens, *40*; ruficaule, *40*; saproxylophilum, 35+; secundum, *40*; subpinnatum, 35+, 338; subsimplex, 338; tristiculum, *40*
Semibarbula orientalis, *40*
serpentine rock, seashore lichens of, 167, 186
Serpotortella chenagonii, 136; madagassa, *136*+; marginata, 136+
SHARP, A. J. (review). A. Noguchi, Illustrated Moss Flora of Japan, 251
SHAW, J. (review). R. E. Magill, Flora of Southern Africa. Byophyta. Part 1 Mosses. Fascicle 2 Gigaspermaceae–Bartramiaceae, 380
SHEARD, J. W. *See* MAYRHOFER, H. & J. W. SHEARD, 106
Shenandoah National Park, Virginia, lichen evidence for changes in air pollution, 21
Sigmatella trichochaete, *40*
Singapore, checklist for mosses of, 24
SLACK, N. G., C. RESCHKE, & B. GILMAN. Scorpidium turgescens rediscovered in New York State, 217
SMITH, R. I. L. Bryophyte oases in ablation valleys on Alexander Island, Antarctica, 45*
SNIDER, J. A. (review). H. A. Crum, Illustrated Moss Flora of Arctic North America and Greenland. 2. Sphagnaceae, 251
SNIDER, J. A. & C. DELGADILLO M. Pleuridium aurantiacum (Musci: Ditrichaceae), a previously undescribed species from central Mexico, 181
SNIDER, J. A., S. M. STUDLAR & M. MEDLEY. A list of the bryophytes of Kentucky, 98
Solorina saccata, 322+
South America, new species of Chrysothrix from, 360
SPENCE, J. R. Bryum Hedw. (Bryaceae) in western North America, 73
Sphaerocarpos texanus, 103
Sphaerophorus globosus, 322+
Sphagnum angustifolium, 365; apollinairci, 132; beccari, *40*; capillifolium, 102; capillifolium var. tenerum, 102; compactum, 102; cuspidatulum, 25+; cuspidatulum var. trengganuense, *40*; cuspidatum, 26; cuspidatum var. flaccidifolium, 26; cuspidatum var. malaccense, *40*; cuspidatum, subsp. subrecurvum, 26; erythrocalyx, 132; fimbriatum, 102, 365; flaccidifolium, *40*; flexuosum, 364+; flexuosum var. **ramosissimum**, var. nov., 364+; fuscum, 104; holttumii, *40*; imbricatum, 102; imbricatum subsp. austinii var. arcticum, *137*; junghuhnianum 26; kelantanense, *40*; limbatum, 132; magellanicum, 40, 102, 335, 365; malaccense, *40*; molle, 102, 104; palustre, 102, 365; pauciporosum, *40*; perichaetiale, 26+; recurvum, 102, 104, 364; roseotinctum, *40*; **rubroflexuosum**, sp. nov., 364+, 365*; russowii, 102; sericeum, 26; sparsum, 335; **steerei**, nom. et stat. nov., 137; strictum, 104; subsecundum, 335, 365; subsecundum var. rufescens, 102; subsecundum var. subsecundum, 102
Sphinctrina tubaeformis, 14+; turbinata, 14+, 322+
Spilonema revertens, 171+, 189
Splachnobryum oorschotii, 30
Splachnum ampullaceum, 161+, mapped, 163; circumboreal species of, 161; luteum, 161+, mapped, 163; rubrum, 161+, mapped, 163; sphaericum, 161+, mapped, 163; vasculosum, 161+, mapped, 164
spores, of Fossombronia, 194*, 195*, 198*, 199*, 200*
Squamidium leucotrichum, 337; livens, 337; nigricans, 337
Staurothele clopima, 88; discedens, redisovered in Alaska, 86+, 87*, 88*, mapped, 89; fissa, 322+; hymenogonia, 88; rufa, 171+, 189; sessilis, 88; sp., 47+; succedens, 88
Stenocybe major, 15+, 322+; pullatula, 15+, 322+
Stenodictyon bisodalense, new to Colombia, 132
Stereocaulon dactylophyllum, 322+; dactylophyllum var. occidentale, 322+, glaucescens, 322+; intermedium, 322+; pileatum, 322+; saxatile, 322+; subcoralloides, 322+; tomentosum, 322+
Stereodon malayanus, *40*
Sterophyllum tavoyense, 34+
Stichococcus mirabilis, 88
Sticta fuliginosa, 322+
Stigmidium marinum, 171+, 189, 322+
Stokesiella praelonga, 102, 179
STOTLER, R. E. & B. CRANDALL-STOTLER. Recent literature on hepatics and anthocerotes—113, 56;—114, 138;—115, 236;—116, 370
Strangospora moriformis, 322+; ochrophora, 322+

STREIMANN, H. The moss genus Papillaria (Meteoriaceae) in New Guinea, 341
Streptopogon calymperes, 336
STUDLAR, S. M. *See* SNIDER, J. A., S. M. STUDLAR & M. MEDLEY, 98
sulfur accumulation, in Flavoparmelia baltimorensis, 21+
sulfur dioxide, effects on lichen communities in Indianapolis, 223
Symphysodon neckeroides, 32
Symphysodontella cylindracea, 32
Syrrhopodon, 18+; aberrans, *40*; aberrans var. laevis, *40*; albidus, 30+; albovaginatus, 30; aristifolius, 19, 30+; borneense, *40*; candidus, *40*; cavifolius, *40*; chenagonii, *136*; ciliatus, 30; ciliatus fo. pseudopodianus, 30; confertus, 30; croceus, 30+; elimbatus, *40*; fallax, *40*; fasciculatus, *40*; fimbriatulus, 30+; flavus, *40*; fraserianus, *40*; gardneri, 30; gaudichaudii, 335; griffithii, *40*; horridulus, *40*; involutus, 30+; japonicus, 19, 30+; laevis, 30+; larminatii, *40*; leprieurii, 335; ligulifolius, *40*; loreus, 30+; louisiadum, *40*; lycopodioides, 335; mammillosus, 51*; manii, *40*; muelleri, 30+; nymanii, *40*; parasiticus, 335; perakensis, *40*; prolifer, 335; pseudo-involutus, *40*; pungens, *40*; quintasii, 55; rectifolius, *40*; repens, *40*; revolutus, *40*; ridleyi, *40*; rigidus, 335; rufescens, *40*; schiffnerianus, *40*; spiculosus, 30+; spiculosus subsp. elimbatus, *40*; spiculosus subsp. herzogii, *40*; spiculosus subsp. horridulus, *40*; subulatus, *40*; texanus, 102; tjibodensis, *40*; trachyphyllus, 30+, 51; treubii, *40*; tristichus, 30; undulatulus, *40*; wallisii, *40*

TAN, B. C. (review). T. Pócs, T. Simon, Z. Tuba & J. Podani (eds.). Proceedings of the IAB Conference of Bryoecology, 158
TAN, B. C. *See* MOHAMED, H. & B. C. TAN, 24
Taxiphyllum deplanatum, 102; taxirameum 37, 102, new to Colombia, 132
Taxithelium acanthocladioides, *40*; alare, 36; albifolium, *40*; bilobatum, *40*; capillipes, 36; decrescens, *40*; deningeri, 36; gottscheanum, *40*; instratum, 36+; isocladioides, 36; isocladum, 36+; kerianum, 36+; lindbergii, 40; magnum var. majus, 36; nepalense, *40*; papillatum, *40*; papillatum var. angustum, *41*; papillatum var. brevifolium, *41*; perakense, *41*; planum, 36+, 339; plumularia, *41*; ridleyi, *41*; singapurense, *41*; stigmosum, *41*; subtrachaelophyllum, *41*; trachaelophyllum, *41*; turgidellum, *41*; vernieri, 36+
Tayloria scabriseta, 132
Tetraphis geniculata, 358; pellucida, 102
Tetraplodon, circumboreal species of, 161; angustatus, 161+, mapped, 164; mnioides, 161+, mapped, 164; urceolatus, 161+, mapped, 164
Tetrodontium brownianum, 358
Thamniopsis killipii, 132, 338; pendula, 338
Thamnium ellipticum, *41*
Thamnobryum alleghaniense, 102; ellipiticum, 33+
Thelia asprella, 102; hirtella, 102; lescurii, 102
Thelotrema lepadinum, 323+
THOMSON, J. W. (review). R. Moberg & E. S. Hansen, The Lichen Genus Physcia and Allied Genera in Greenland, 69
THOMSON, J. W. & B. M. MURRAY. Staurothele discedens and Gyalidea lecideopsis var. convarians rediscovered in Alaska, together with Polyblastia cucurbitula, sp. nov., 86
THOR, G. Three calicialian lichens new to the continental United States, 367
THOR, G. Two new species of Chrysothrix from South America, 360
Thrombium epigaeum, 323+
Thuidium abietinum, 102; acuminatum, *120*; antillarum, 116+, 132, 338; bifarium, *41*; cymbifolium, 34; delicatulum, 93, 102; glaucinoides, 34; glaucinum, 34; leptocladum, 132; meyenianum, 34; minutulum, 102, 104, 132; orientale, 34; peruvianum, 338; plumulosum, 34; pygmaeum, 102; recognitum, 102; scitum, 104; tamariscinum, growth rate of, 219; trachypodum, *41*; urceolatum, 116+, 132, 338
Thyridium fasciculatum, *41*; flavum, *41*; manii, *40*; repens, *41*; undulatulum, *41*; undulatum, *41*
Thysanomitrium abbreviatum, *41*; exasperatum, *41*; ridleyi, *41*; umbellatum, *41*
TIMME, S. L. & F. D. BOWERS. Pleurochaete squarrosa (Brid.) Lindb. new to Mississippi and extensions of its range in Alabama, 368
Timmia megapolitana, 102
Tomenthypnum nitens, effects of simulated acid rain on, 121
Tortella fragilis var. tortelloides, 47+; humilis, 102; princeps, 47+
Tortula andicola, 19, 132; chisosa, 55; intermedia, 18; mniadelphus, new to Colombia, 132; mniifolia, 336; pagorum, 102; papillosa, 102; ruralis, 18
Trachyloma indicum, 32
Trachypus bicolor, 337; humilis, 32
Trachythecium calciçolum, 37; verrucosum, 37+
Trapelia coarctata, 323+; involuta, 323+; placodioides, 323+
Trapeliopsis flexuosa, 323+; granulosa, 323+; viridescens, 323+
Trematodon acutus, *41*; longicollis, 28, 102; montanus, 357
Trichocolea tomentella, 103
Trichosteleum albifolium, 36; boschii, 36, 40; brachypelma, 36; brachypelma var. compactum, 36; bruchii, *41*; cyparissoides, 339; densirameum, *41*; fruticolum, 36; hamatum, 36; laxirete, *41*; leptocarpon, *41*; luxurians, *41*; mammosum, 36; monostictum, 36; papillatum, *41*; papillosum, 339; pinnatum, 36; procerum, 36; pseudoinstratum, *41*; schlimii, 339; singapurense, 36
Trichostomum brachydontium, 336; orientale, *41*; sarawakense, 30; tenuirostre, 102
Trismegistia brauniana, 36+; calderensis, 36; complanatula, 36; lancifolia, 36; lancifolia var. korthalsii, 36; rigida, 36
Tristichella spiculifera, 36
Tritomaria exsecta, 103
Tuckermannopsis americana, 323+; ciliaris, 323+; halei, *323+*; orbata, 323+
Tylophoron protrudens, new to the United States, 367

Uleobryum curtisii, 214+
Ulota coarctata, 102; crispa, 102; drummondii, 358; hutchinsiae, 102
Umbilicaria decussata, 47+; deusta, 323+; mammulata, 323+; muehlenbergii, 323+; papulosa, *323+*; vellea, 323+
Usnea antarctica, 47+; filipendula, 323+; fulvoreagens, *323+*; hirta, 323+; lapponica, 323+; longissima, 323+; rubicunda, new to Canada, 323+; sphacelata, 47+, 108; strigosa, 323+; subfloridana, 323+; trichodea, 323+

Verrucaria aethiobola, 323+; amphibia, 189; ceuthocarpa, 171+, 189, 323+; degelii, 171+, 186+, 323+; discedens, *86*; ditmarsica, 323+, durietzii, 189; elaeomelaena, 189; erichsenii, 171+, 189, 323+; halizoa, 171+, 189, 323+; laevata var. nigrita, 189; lobata,

189; maura, 171+, 188+, 323+; microspora, *171*, 323+; mucosa, 171+, 189, 323+; muralis, 323+; nigrescens, 189, 323+; nigrescentoidea, 323+; sandstedei, 171+, new to North America, 186+; spp., 171+, 189; striatula, 171+, 189, 323+
Vesicularia amphibola, 339; dubyana, 37; kurzii, 37; miquelii, 37+; montagnei, 37; muellerianum, 41; reticulata, 37+; vesicularis, new to Colombia, 339
Virginia, lichen evidence for changes in air pollution in Shenandoah National Park, 21
VITT, D. H. *See* ROCHEFORT, L. & D. H. VITT, 121

Warburgiella complanata, 36; leptocarpa, 36+; pycnophylla, 36+
Washington, seashore lichens of Fidalgo Island, 167, 186
Webera elongata, *41*
Weissia controversa, 102, rhizoid gemmae in, 184*+; edentula, 30+; malayensis, 30+; sharpii, 102
WHITTEMORE, A. (review). B. N. McKnight, The Bryophytes of Illinois: An Annotated and Indexed Bibliography and Checklist, 253
Wijkia alboalaris, 36+; hornschuchii, 36; polymorpha, 36+; surcularis, *41*; tanytricha, 36+; tenuisetum, 41
Wilsoniella acutifolia, *41*; decipiens, 28+; decipiens var. acutifolia, 28+; pellucida, *41*

Xanthoparmelia centrifuga, *323*+; conspersa, 323+; cumberlandia, 323+; taractica, 323+
Xanthoria candelaria, 171+; elegans, 47+; fallax, 226, 324+; hasseana, *324*+; parietina, 178, 324+; polycarpa, 324+; sorediata, 324+; sp. affin. elegans, 324+
Xylographa abietina, 324+; opegraphella, 324+

ZANDER, R. H. *See* REESE, W. D. & R. H. ZANDER, 18, 136
Zelometeorium patulum, 337; recurvifolium, 337